

STIC-ILL

From: Murphy, Joseph
Sent: Monday, November 29, 1999 3:36 PM
To: STIC-ILL
Subject: 08920272

1646

272588

272768

Please Rush me the following references:

Calof AL, et al

Regulation of neurogenesis and neuronal differentiation in primary and immortalized cells from mouse olfactory epithelium.
Ciba Found Symp. 1991;160:249-65; discussion 265-76.
PMID: 1752166; UI: 92090184.

ZGE-991130845

Pixley SK, et al.

Cultured rat olfactory neurons are excitable and respond to odors.
Brain Res Dev Brain Res. 1990 Apr 1;53(1):125-30.
PMID: 2350878; UI: 90275756.

Pixley SK.

Characterization of olfactory receptor neurons and other cell types in dissociated rat olfactory cell cultures.
Int J Dev Neurosci. 1996 Nov;14(7-8):823-39.
PMID: 9010728; UI: 97163981.

Noble M, et al.

The growth of olfactory neurons in short-term cultures of rat olfactory epithelium.
Neurosci Lett. 1984 Mar 23;45(2):193-8.
PMID: 6728313; UI: 84220439.

Maue RA, et al.

Preparation of isolated mouse olfactory receptor neurons.
Pflugers Arch. 1987 Jul;409(3):244-50.
PMID: 3627945; UI: 87316795.

V. NO
11/30

Schubert D, et al.

Ontogeny of electrically excitable cells in cultured olfactory epithelium.
Proc Natl Acad Sci U S A. 1985 Nov;82(22):7782-6.
PMID: 2866512; UI: 86067989.

Wolozin B, et al.

Continuous culture of neuronal cells from adult human olfactory epithelium.
J Mol Neurosci. 1992;3(3):137-46.
PMID: 1320921; UI: 92329300.

Sosnowski JS, et al.

Chemical traumatization of adult mouse olfactory epithelium in situ stimulates growth and differentiation of olfactory neurons in vitro.
Brain Res. 1995 Dec 8;702(1-2):37-48.
PMID: 8846094; UI: 96255746.

Eustache I, et al.

Effects of insulin-like growth factors on organotypic cocultures of embryonic rat brainstem slices and skeletal muscle fibers.
Brain Res Dev Brain Res. 1994 Sep 16;81(2):284-92.
PMID: 7813048; UI: 95112382.

NSL 02631

THE GROWTH OF OLFACTORY NEURONS IN SHORT-TERM CULTURES OF RAT OLFACTORY EPITHELIUM

MARK NOBLE¹, PETER S. MALLABURN² and NIGEL KLEIN²

¹Department of Clinical Neurology, Institute of Neurology, Queen Square, London WC1N 3BG and

²Department of Anatomy and Embryology and Centre for Neuroscience, University College London, Gower Street, London WC1E 6BT (U.K.)

(Received November 4th, 1983; Revised version received January 9th, 1984; Accepted January 10th, 1984)

Key words: olfactory neuron – culture – astrocytes – neurogenesis – rat

We have found that purified astrocytes will support the growth of olfactory epithelial neurons (OENs) *in vitro*. Candidate OENs were visible by light microscopy within one day after plating of dissociated cells from neonatal rat ethmoturbinates. Electron microscopy showed that OENs expressed the unique ultrastructural features which characterize these cells *in situ*, showing that regulation of neuronal morphology and ultrastructure can function independently of contact with other cells of the appropriate tissue. Characteristics of the cultures are such that this *in vitro* system would be useful in electrophysiological investigations of the chemosensory olfactory neurons.

The primary chemosensory neurons of the olfactory epithelium are the only population of neurons continuously replaced throughout adult life. Complete cycles of development, beginning with dividing neuronal precursors and continuing through neuronal maturation to cell death, occur over periods of 3–4 weeks [5, 11]. These cycles of loss and replacement mean that denervation and appropriate reinnervation of the olfactory bulb must constantly be recurring [11]. Thus, the olfactory neuroepithelium is a useful model system for studying several major aspects of neuronal development and synaptogenesis. Olfactory neurons are also of particular physiological interest, because the mechanism by which they discriminate different odors is unknown; it is not yet even established whether single neurons respond to single odors or instead recognize a variety of odor stimuli.

A major obstacle to the usefulness of the olfactory epithelium in studies of neurogenesis, or of the electrophysiology of chemosensory transduction, has been the inability to grow olfactory epithelial neurons (OENs) in tissue culture as single cells, free of supporting sustentacular cells, other neurons and the nasal mucosa. As a first step towards investigating the regulation of OEN development, we have developed techniques for growing these cells *in vitro*. In this paper we describe the growth and ultrastructure of OEN grown on monolayers of purified astrocytes, a culture system in which these neurons can be identified within one day after plating and are subsequently accessible for physiological, morphological and immunochemical studies. In addition, we have found that even when removed from their normal site of growth and complement of supporting cells, OENs express the unique ultrastructural features which characterize these neurons *in situ*.

ion, and that the recorded
stant recordings from the

n the dendrites usually have
an the responses generated
around 100 ms, while in the
hundred milliseconds. It is
ings in that study were from
iV, i.e. within the potential
om Purkinje cell somata.
e soma is, it is possible that
sponses in depolarized den-
nt results do not cast doubt
e can also be seen as unitary
field potentials from the

Faculty, University of Lund
ical Research Council (Pro-

1 cerebellar cortex of the cat, J.

dendritic plateau potentials evoked
1, 340 (1983) 209–223.

tween responses in Purkinje cells
cat, J. Physiol. (Lond.) 340 (1983)

ig fibre responses recorded from

neural Machine, Springer-Verlag,

ited in Purkinje cell dendrites by
1) 207–221.

bing fibre input to Purkinje cells:
alay (Eds.), The Cerebellum: New

cerebellar Purkinje cell responses

Purified cortical astrocytes were prepared by a modification [12] of methods described previously [9]. Purified astrocytes were passaged once in tissue culture flasks, and then grown as monolayers on poly-L-lysine coated glass coverslips. Astrocytes obtained in this way produced flat monolayers, with good visualization of added neurons. Greater than 95% of the cells labeled with rabbit anti-glial fibrillary acidic protein antiserum (a specific marker for astrocytes [1, 13]), and there were no neurofilament⁺ cells (i.e. neurons [15]), galactocerebroside⁺ cells (i.e. oligodendrocytes [13, 14]) or normal rabbit serum-binding cells (i.e. macrophages/microglia [13]). The non-astrocytes had fibroblast-like morphologies and often expressed cell-surface fibronectin.

The nasal ethmoturbinates were dissected from newborn Sprague-Dawley rats (N. Klein et al., in preparation) and mechanically dissociated into single cell suspensions [6]. The yield of cells per animal obtained was low ($\approx 2.5 \times 10^5$ /rat), but mechanical dissociation offered the major advantage of allowing easy separation of associated cartilage by passage through 45- μ m mesh nylon gauze. Cells were diluted to 2, 5 or 10×10^4 cells/ml in Dulbecco's Modified Eagle's Medium (DMEM) containing 5% horse serum, 5% fetal calf serum, 2 mM glutamine and 25 μ g/ml gentamicin. Cell suspensions were plated in 0.5 ml aliquots onto astrocyte monolayers growing on glass coverslips in Falcon Multiwell Trays (No. 3047). Cultures were grown for 48–72 h before fixation and preparation for electron microscopy (see Figs. 1 and 2).

Within 24 h after plating, many cells in the cultures possessed morphological features in common with OENs in situ. Candidate OENs were bipolar, with one short process ($< 30 \mu$ m) terminating in a bulbous expansion and a longer ($> 250 \mu$ m) and thinner process, each originating from opposite poles of an elliptical (ca. $17 \times 6 \mu$ m) cell body. The number of such cells per culture was usually 1–2% of the total number of neuroepithelial cells originally plated. No bipolar cells were found in cultures of bronchial respiratory epithelium, the most likely contaminating tissue in this preparation, indicating that these cells were probably derived from olfactory neuroepithelium. To confirm this, an ultrastructural comparison was performed between candidate OENs and OENs of the newborn rat ethmoturbinate.

Mature OENs in situ display a stereotyped and distinctive ultrastructure [5] (Fig. 1). The major distinguishing feature is the olfactory club (Fig. 1a), which is located at the tip of the apical dendrite and projects into the olfactory mucosa. The club contains numerous centrioles/basal bodies, and often possesses ciliary rootlets and cilia. The dendrite itself is usually 15–30 μ m long and 1–3 μ m in diameter and contains numerous microtubules and long slender mitochondria aligned along its long axis (Fig. 1b). Centrioles/basal bodies are often seen in the dendrites of less mature neurons, presumably in transit to the sensory club. The dendrites contain no 10-nm neurofilaments. The elliptical cell body is centrally placed in the epithelium and typically measures $17 \times 6 \mu$ m (Fig. 1c). The cytoplasm contains abundant free ribosomes, with a sparse polar distribution of other organelles, including rough endoplasmic reticulum (rER) and mitochondria with Golgi apparatus evident proximal to processes in younger cells. The nucleus is ovoid, with large clumps of chromatin

a modification [12] of methods passed once in tissue culture lysine coated glass coverslips. monolayers, with good visualization is labeled with rabbit anti-glial ker for astrocytes [1, 13]), and [15]), galactocerebroside⁺ cells bbit serum-binding cells (i.e. ad fibroblast-like morphologies

newborn Sprague-Dawley rats dissociated into single cell suspension was low ($\approx 2.5 \times 10^5$ /rat), but ge of allowing easy separation of a nylon gauze. Cells were diluted l Eagle's Medium (DMEM) con- M glutamine and 25 μ g/ml gen- quots onto astrocyte monolayers trays (No. 3047). Cultures were on for electron microscopy (see

ultures possessed morphological e OENs were bipolar, with one expansion and a longer (> 250 opposite poles of an elliptical (ca. per culture was usually 1-2% of y plated. No bipolar cells were m, the most likely contaminating ells were probably derived from a structural comparison was per- ie newborn rat ethmoturbinate. distinctive ultrastructure [5] (Fig. y club (Fig. 1a), which is located the olfactory mucosa. The club ten possesses ciliary rootlets and and 1-3 μ m in diameter and con- ochondria aligned along its long n in the dendrites of less mature The dendrites contain no 10-nm ly placed in the epithelium and oplasm contains abundant free r organelles, including rough en- olgi apparatus evident proximal with large clumps of chromatin




Fig. 1. The figure shows the fine structure of olfactory neurons in situ. Tissue was processed for electron microscopy by immersion prefixation in 2.5% glutaraldehyde for 2 h followed by post-fixation in 2.0% OsO₄, also for 2 h. Both fixatives were buffered in PBS (Dulbecco 'A'). After dehydration in ethanol, specimens were embedded in Araldite. a: two sensory clubs are shown projecting from the free surface of the epithelium. Abundant centrioles are seen (Ce) both in, and en route to, the club. Both clubs make extensive contact (white triangles) with supporting cells (S Cell). Bar = 0.5 μ m. b: a low power micrograph showing a dendrite whose sensory club (SC) has not yet penetrated the free surface, as confirmed by serial section analysis (not shown). Elongate mitochondria (M) and microtubules (Mt) are orientated along the dendrite's long axis. Bar = 1.5 μ m. c: detail of an olfactory cell body orientated as in (b) and located proximal to the dendrite. Note the single nucleus (N) and the characteristically distributed chromatin (C) located in the nucleus. Sparse organelles, especially rough endoplasmic reticulum (rER) and mitochondria (M), are seen in a polar cytoplasm, with profuse free ribosomes distributed throughout. Bar = 2 μ m.

and a single nucleolus distributed in an electronlucent nucleoplasm. A slender axon (0.3-0.5 μ m) projects through the basement membrane and lamina propria of the epithelium and eventually to the olfactory bulb.

In vitro, candidate cells expressed the characteristic ultrastructural features of OENs in situ (Fig. 2). A typical pear-shaped bipolar cell is shown in Fig. 2a, with one long slender process (which extends > 250 μ m) and a thicker short ($\approx 25 \mu$ m) process ending in a bulbous expansion; Fig. 2b is an electron microscopic montage of the same cell. The 25- μ m process contains numerous aligned microtubules, elongated mitochondria and at least one centriole (Fig. 2c). The most important feature is the club-shaped specialization terminating the 25- μ m process (Fig. 2d). This club contains less than 25 centrioles and at least one ciliary rootlet; segregation of centrioles to such a specialization is a unique feature of OENs. The elliptical cell body contains rER profiles and mitochondria, especially proximal to the long process, and profuse free ribosomes. The nucleus possesses the chromatin pattern characteristic of OENs in situ.

Twelve of 15 candidate OENs chosen at the light microscopic level had the defining ultrastructural features of OENs. Two of the negative cells had some, but not




Fig. 2. The figure shows the morphology and fine structure of olfactory neurons in vitro. Cultures were processed for electron microscopy as above, except that the times of each fixation were reduced to 15 min. Cultures and coverslips were embedded in a thin layer of Araldite to facilitate light microscopic study. a: a light micrograph of a candidate OEN lying on an astrocyte. This was photographed in Araldite, then serially sectioned en face for electron microscopy, the coverslip having been removed with hydrofluoric acid. Bar = 10 μ m. b: a through-section photomontage of the candidate OEN shown in (a). Note the characteristic elliptical cell body (CB) and nucleus, the dendrite (D) projecting from one pole, and the segregated centrioles in a terminal sensory club (SC). A long, slender axon (A) projects from the opposite pole. The smaller cell may be an immature neuron. Bar = 5 μ m. c: detail of the dendrite shows that abundant microtubules (Mt) are present together with slender, elongate mitochondria (M) and a solitary centriole (Ce). A point of contact with the smaller cell is shown (large arrow), as is the astrocyte on which the OEN rests (A). Bar = 0.5 μ m. d: detail of the sensory club. The accumulation of centrioles (Ce) is obvious; also present is a ciliary rootlet (Cr). The two filopodial processes and the large inclusion body are culture artefacts. Bar = 0.5 μ m.

all of the features of OENs, and the remaining one cell was indistinguishable from a fibroblast. Various minor features poorly seen in the cell in Fig. 2, and associated




Figure 2 shows an olfactory neuron in vitro. Cultures were fixed at various times of each fixation were reduced to 15 minutes of Araldite to facilitate light microscopic examination of an astrocyte. This was photographed in situ, the coverslip having been removed with a montage of the candidate OEN shown in Figure 1c. The dendrite (D) projecting from one side of the cell body (SC). A long, slender axon (A) projects from the other side of the cell body. Bar = 5 μ m. c: detail of the dendrite with slender, elongate mitochondria (M). A smaller cell is shown (large arrow), as is the tail of the sensory club. The accumulation of filopodia (Cr). The two filopodial processes and the

ne cell was indistinguishable from the cell in Fig. 2, and associated

with differing stages of OEN development in situ, were often found in candidate OENs (Fig. 3), suggesting that cultures contained OENs at various stages of maturation. In addition, cells which did not wholly conform to our light microscopic criteria for candidate OENs often displayed various ultrastructural features of OENs. Thus, it is likely that we are underestimating the overall frequency of OENs in our cultures.

The ability to support neuronal growth in vitro is one of the distinguishing characteristics of the astrocyte, and astrocytes have been found to promote the growth of various PNS and CNS neurons [8, 12]. We have now found that astrocytes will support the in vitro growth of OENs, a population of neurons which has proven difficult to grow in tissue culture conditions.

Of particular interest was the observation that OENs could be readily identified by their expression in vitro of ultrastructural features unique to this cell type. It has been observed that some classes of neurons will develop morphologies in vitro similar to those seen in situ [7]. Our findings appear to offer the first demonstrations that the regulation of morphological development can extend to the level of ultrastructure and that this regulation will function independently of contact with other cells of the appropriate tissue.

Will normal cycles of OEN development occur in vitro, with associated cell division, neuronal maturation and neuronal death? One previous study, in which antiserum against olfactory marker protein (OMP) was used to identify OENs, suggested that in vitro development did not occur [4]. However, it may be that, as with Schwann cells and myelin-specific compounds [2, 10], OMP production requires induction by specific signals, while generation of normal morphology does not. More recent studies [3, 6], in which cultures of olfactory epithelial tissue were followed




Fig. 3. This figure shows other features of olfactory neurons in vitro. a: an OEN cell body, orientated as in Fig. 1c, showing an extensive Golgi system (G) proximal to the dendrite and club (not shown). Such Golgi systems are often seen in association with growing processes, and its absence in Fig. 1c and 2b suggests that the latter are mature neurons. Bar = 1.5 μ m. b: a high power micrograph of part of a sensory club showing elongated mitochondria (M) and microtubules (Mt), with many centrioles (Ce). A group of 8 orientated centrioles (OCe) demonstrate a regularity often seen in situ.

for up to 2 months, suggest that neurogenesis may well occur in vitro under appropriate conditions; our preliminary work on long-term cultures also suggests that neurogenesis occurs in the system reported here. Advantages to growth on astrocyte monolayers are that many OEN cultures can be obtained from 2-3 newborn rats, and OENs grow as single cells easily recognizable by light microscopy within a short time after plating. Thus, this culture system will be useful in selecting antibodies which recognize antigens on the OEN cell surface, and is eminently suitable for electrophysiological studies of odor sensitivities and of chemosensory transduction.

It is a pleasure to acknowledge the advice and support of Martin Raff, Bob Lieberman and Eva Franke, and also of Duncan O'Dell, Derek James, Derek Fraser and Paul Debbage. M.N. is supported by an MRC Project Grant.

- 1 Bignami, A., Eng, L.F., Dahl, D. and Uyeda, C.T., Localization of the glial fibrillary acidic protein in astrocytes by immunofluorescence, *Brain Res.*, 43 (1972) 429-435.
- 2 Brookes, J.P., Fields, K.L. and Raff, M.C., Studies on cultured rat Schwann cells. I. Establishment of purified population from cultures of peripheral nerve, *Brain Res.*, 165 (1979) 105-118.
- 3 Debbage, P.L., Klein, N.J., O'Dell, D.S., Fraser, D.A. and James, D.W., The culture of olfactory neurons, *J. Anat.*, 135 (1982) 816-817.
- 4 Farbman, A.I. and Margolis, F.L., Olfactory marker protein during ontogeny: immunohistochemical localization, *Develop. Biol.*, 74 (1980) 205-215.
- 5 Graziadei, P.P.C. and Monti-Graziadei, G.A., Neurogenesis and neuron regeneration in the olfactory system of mammals. I. Morphological aspects of differentiation and structural organization of olfactory sensory neurons, *J. Neurocytol.*, 8 (1979) 1-18.
- 6 Klein, N.J., Debbage, P.L., Fraser, D.A., Mallaburn, P.S., O'Dell, D.S. and James, D.W., Rat olfactory epithelium in vitro and in situ: a comparison, *Europ. J. Cell Biol., Suppl.* 1 (1983) 24.
- 7 Kriegstein, A.R. and Dichter, M.A., Morphological classification of rat cortical neurons in cell culture, *J. Neurosci.*, 3 (1983) 1634-1647.
- 8 Lindsay, R.M., Adult rat brain astrocyte support survival of both NGF-dependent and NGF-insensitive neurons, *Nature (Lond.)*, 282 (1980) 80-82.
- 9 McCarthy, K.D. and de Vellis, J., Preparation of separate astroglial and oligodendroglial cultures from rat cerebral tissue, *J. Cell Biol.*, 85 (1980) 890-902.
- 10 Mirsky, R., Winter, J., Abney, E.R., Pruss, R.M., Gavrilovic J. and Raff, M.C., Myelin-specific proteins and glycolipids in rat Schwann cells and oligodendrocytes in culture, *J. Cell Biol.*, 84 (1980) 483-494.
- 11 Monti-Graziadei, G.A. and Graziadei, P.P.C., Neurogenesis and neuron regeneration in the olfactory system of mammals. II. Degeneration and reconstitution of the olfactory sensory neurons after axotomy, *J. Neurocytol.*, 8 (1979) 197-213.
- 12 Noble, M., Fok-Seang, J. and Cohen, J., Glia are a unique substrate for the in vitro growth of CNS neurons, submitted.
- 13 Raff, M.C., Fields, K.L., Hakomori, S., Mirsky, R., Pruss, R.M. and Winter, J., Cell-type-specific markers for distinguishing and studying neurons and the major classes of glial cells in culture, *Brain Res.*, 174 (1979) 283-308.
- 14 Raff, M.C., Mirsky, R., Fields, K.L., Lisak, R.P., Dorfman, S.H., Silberberg, D.H., Gregson, N.A., Liebowitz, S. and Kennedy, M., Galactocerebroside: a specific cell surface antigenic marker for oligodendrocytes in culture, *Nature (Lond.)*, 274 (1978) 813-816.
- 15 Wood, J.D. and Anderton, B., Monoclonal antibodies to mammalian neurofilaments, *Bioscience Rep.*, 1 (1981) 263-268.